

### V 主要病害虫発生程度別面積及び防除面積

作物名	病害虫名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
水稲 103,000 ha	葉いもち	93	0.1	0	0.0	102,907	93	0	0	0	102,079	249,355	やや遅	少
	穂いもち	94	0.1	6	0.0	102,906	88	0	6	0	102,079	249,355	やや遅	少
	紋枯病	21,223	20.6	4,960	4.8	81,777	16,263	3,749	1,211	0	11,040	14,164	並	多
	縞葉枯病	939	0.9	0	0.0	102,061	939	0	0	0	66,598	98,819	—	並
	ばか苗病	1,139	1.1	0	0.0	101,861	1,139	0	0	0	99,468	107,426	—	並
	苗立枯病	1,495	1.5	0	0.0	101,505	1,495	0	0	0	87,855	90,529	—	並
	種子伝染性細菌病	98	0.1	14	0.0	102,902	84	14	0	0	98,784	98,784	—	少
	ニカメイガ	582	0.6	0	0.0	102,418	582	0	0	0	54,443	95,606	やや早	やや少
	セジロウンカ	2,336	2.3	0	0.0	100,664	2,336	0	0	0	98,384	207,328	早	やや多
	ヒメトビウンカ	17,780	17.3	146	0.1	85,220	17,634	146	0	0	97,703	207,162	早	並
	イネドロオイムシ	5,481	5.3	6	0.0	97,519	5,475	6	0	0	97,722	110,589	やや早	少
	アカヒゲ ホソミドリ カスミカメ	47,058	45.7	101	0.1	55,942	46,957	101	0	0	102,586	249,808	やや早	やや少
	イネミギワバエ	2,489	2.4	641	0.6	100,511	1,848	641	0	0	13,523	15,416	並	並
フタオビコヤガ	2,346	2.3	15	0.0	100,654	2,331	15	0	0	78,636	101,291	並	少	
秋まき小麦 103,768 ha	赤さび病	19,063	18.4	3,098	3.0	84,705	15,965	2,788	297	13	103,768	242,647	やや早	やや多
	うどんこ病	4,921	4.7	77	0.1	98,847	4,844	77	0	0	103,438	254,209	やや早	やや少
	雪腐病類	12,529	12.1	1,466	1.4	91,239	11,063	1,273	152	41	95,255	95,255	—	少
	眼紋病	4,166	4.0	281	0.3	99,602	3,885	281	0	0	20,010	20,010	—	やや少
	赤かび病	22,436	21.6	3,233	3.1	81,332	19,203	3,233	0	0	103,768	355,072	—	やや少
春まき小麦 (春まき) 16,594 ha	赤かび病	3,023	18.2	522	3.1	13,571	2,501	518	4	0	16,594	57,881	—	並
	ムギキモグリバエ	944	5.7	0	0.0	15,650	944	0	0	0	6,086	10,198	やや早	並
春まき小麦 (初冬まき) 1,038 ha	赤かび病	568	54.7	7	0.7	470	561	4	3	0	1,038	4,119	—	やや少
	ムギキモグリバエ	22	2.1	0	0.0	1,016	22	0	0	0	137	137	やや早	やや少

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積(ha)	率(%)	面積(ha)	率(%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
大豆 39,100 ha	べと病	8,034	20.5	182	0.5	31,066	7,852	172	10	0	19,792	23,664	並	やや少
	わい化病	4,907	12.5	7	0.0	34,193	4,900	7	0	0	39,056	53,018	—	並
	マメシクイガ	5,821	14.9	192	0.5	33,279	5,629	192	0	0	39,005	83,018	並	少
	食葉性 鱗翅目幼虫	11,403	29.2	316	0.8	27,697	11,087	313	3	0	32,683	56,253	並	並
	タネバエ	365	0.9	0	0.0	38,735	365	0	0	0	39,100	39,100	—	並
小豆 20,900 ha	菌核病	2,074	9.9	9	0.0	18,826	2,065	9	0	0	20,803	60,406	並	やや少
	灰色かび病	2,495	11.9	28	0.1	18,405	2,467	28	0	0	20,844	60,129	並	やや少
	茎疫病	463	2.2	34	0.2	20,437	429	34	0	0	3,529	4,805	—	少
	落葉病	380	1.8	0	0.0	20,520	380	0	0	0	0	0	—	少
	食葉性 鱗翅目幼虫	3,423	16.4	0	0.0	17,477	3,423	0	0	0	20,284	25,959	並	少
菜豆 6,340 ha	菌核病	1,261	19.9	21	0.3	5,079	1,240	21	0	0	6,182	20,000	やや早	並
	灰色かび病	1,034	16.3	4	0.1	5,306	1,030	4	0	0	6,182	20,000	並	並
	黄化病	105	1.7	16	0.3	6,235	89	16	0	0	6,340	8,547	—	並
	タネバエ	16	0.3	0	0.0	6,324	16	0	0	0	6,340	6,340	—	少
ばれいしょ 49,600 ha	疫病	3,785	7.6	719	1.4	45,815	3,066	541	178	0	49,600	371,498	並	少
	塊茎腐敗	2,646	5.3	233	0.5	46,954	2,413	233	0	0	49,600	371,498	—	やや少
	軟腐病	7,355	14.8	82	0.2	42,245	7,273	82	0	0	49,377	170,696	—	少
	黒あし病	444	0.9	0	0.0	49,156	444	0	0	0	39,209	39,209	—	並
	そうか病	9,897	20.0	2,478	5.0	39,703	7,419	1,945	355	178	39,458	39,458	—	並
	粉状そうか病	3,089	6.2	30	0.1	46,511	3,059	30	0	0	5,090	5,090	—	並
	アブラムシ類	4,790	9.7	64	0.1	44,810	4,726	51	13	0	49,600	152,184	やや早	少
てんさい 56,700 ha	褐斑病	6,782	12.0	499	0.9	49,918	6,283	404	95	0	56,700	304,888	並	少
	根腐病 (黒根病を含む)	2,145	3.8	3	0.0	54,555	2,142	3	0	0	56,700	111,439	—	少
	そう根病	55	0.1	0	0.0	56,645	55	0	0	0	0	0	—	並
	黄化病 (西部萎黄病)	768	1.4	0	0.0	55,932	768	0	0	0	55,954	188,086	—	少
	ヨトウガ (第1回)	11,183	19.7	1,435	2.5	45,517	9,748	1,435	0	0	56,700	87,263	やや早	やや多
	ヨトウガ (第2回)	6,652	11.7	291	0.5	50,048	6,361	291	0	0	56,700	79,527	やや早	並
	テンサイ モグリハナバエ	70	0.1	0	0.0	56,630	70	0	0	0	37,476	45,595	—	少

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
たまねぎ 14,600 ha	白斑葉枯病	3,783	25.9	267	1.8	10,817	3,516	267	0	0	14,600	101,433	並	少
	軟腐病	6,648	45.5	247	1.7	7,952	6,401	247	0	0	14,600	104,250	—	やや少
	乾腐病	1,404	9.6	24	0.2	13,196	1,380	24	0	0	0	0	—	少
	タマネギバエ、 タネバエ	1,767	12.1	36	0.2	12,833	1,731	36	0	0	5,026	5,466	—	やや多
	ネギアザミウマ	14,013	96.0	2,232	15.3	587	11,781	1,945	283	4	14,600	87,387	やや早	やや多
	ネギハモグリ バエ	1,099	7.5	10	0.1	13,501	1,089	6	4	0	13,777	52,422	並	少
ねぎ 623 ha	さび病	45	7.2	0	0.0	578	45	0	0	0	623	1,696	—	少
	ネギアザミウマ	408	65.5	194	31.1	215	214	23	93	78	623	3,496	早	やや多
にんじん 4,670 ha	黒葉枯病	1,052	22.5	86	1.8	3,618	966	86	0	0	4,469	13,010	—	やや少
だいこん 3,250 ha	軟腐病	1,146	35.3	204	6.3	2,104	942	161	43	0	3,065	9,026	—	並
	キスジ トビハムシ	809	24.9	26	0.8	2,441	783	26	0	0	3,250	5,944	—	やや少
キャベツ 1,170 ha	コナガ	553	47.3	42	3.6	617	511	36	6	0	1,170	6,607	並	やや少
	ヨトウガ	519	44.4	36	3.1	651	483	32	4	0	1,170	6,028	やや早	並
ブロッコリー 2,700 ha	コナガ	1,588	58.8	447	16.6	1,112	1,141	439	8	0	2,700	9,721	並	やや多
	ヨトウガ	1,314	48.7	158	5.9	1,386	1,156	131	27	0	2,700	7,707	やや早	並
りんご 516 ha	モニリア病	5	1.0	0	0.0	511	5	0	0	0	516	972	やや遅	少
	黒星病	229	44.4	0	0.0	287	229	0	0	0	516	4,680	やや早	やや多
	斑点落葉病	88	17.1	0	0.0	428	88	0	0	0	516	2,296	並	並
	腐らん病	435	84.3	114	22.1	81	321	103	11	0	456	1,316	—	多
	ハマキムシ類	78	15.1	0	0.0	438	78	0	0	0	516	2,399	早	並
	モモシンクイガ	81	15.7	0	0.0	435	81	0	0	0	516	2,340	早	やや少
	ハダニ類	209	40.5	3	0.6	307	206	3	0	0	516	1,495	並	やや少
	キンモンホソガ	129	25.0	0	0.0	387	129	0	0	0	516	1,608	並	並
	ネズミ類	40	7.8	0	0.0	476	40	0	0	0	23	23	—	やや少