

R4年 主要病害虫発生程度別面積及び防除面積

作物名	病害虫名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
水稲 96,100 ha	葉いもち	247	0.3	0	0.0	95,853	247	0	0	0	95,501	231,435	並	やや少
	穂いもち	197	0.2	0	0.0	95,903	197	0	0	0	95,501	231,435	やや遅	やや少
	紋枯病	22,975	23.9	6,116	6.4	73,125	16,859	4,076	1,503	537	13,866	16,791	並	やや多
	縞葉枯病	759	0.8	135	0.1	95,341	624	135	0	0	66,435	88,985	-	並
	ばか苗病	1,175	1.2	0	0.0	94,925	1,175	0	0	0	96,100	96,100	-	並
	苗立枯病	1,193	1.2	0	0.0	94,907	1,193	0	0	0	86,605	98,270	-	並
	種子伝染性細菌病	82	0.1	0	0.0	96,018	82	0	0	0	96,100	96,100	-	少
	ニカメイガ	755	0.8	0	0.0	95,345	755	0	0	0	46,404	84,477	並	並
	セジロウンカ	2,476	2.6	0	0.0	93,624	2,476	0	0	0	93,117	197,963	やや早	並
	ヒメトビウンカ	8,761	9.1	252	0.3	87,339	8,509	252	0	0	91,752	193,262	やや早	やや少
	イネドロオイムシ	1,313	1.4	0	0.0	94,787	1,313	0	0	0	91,467	100,039	並	少
	アカヒゲ ホソミドリ カスミカメ	25,646	26.7	477	0.5	70,454	25,169	477	0	0	95,758	224,797	早	並
	イネミギワバエ	284	0.3	0	0.0	95,816	284	0	0	0	28,753	28,753	遅	少
フタオビコヤガ	1,473	1.5	0	0.0	94,627	1,473	0	0	0	58,516	70,262	並	少	
秋まき小麦 108,500 ha	赤さび病	23,885	22.0	1,456	1.3	84,615	22,429	1,456	0	0	108,492	257,052	並	並
	うどんこ病	4,970	4.6	131	0.1	103,530	4,839	131	0	0	108,485	262,216	並	並
	雪腐病類	46,324	42.7	6,320	5.8	62,176	40,004	5,746	467	107	102,102	109,610	-	やや多
	眼紋病	4,623	4.3	576	0.5	103,877	4,047	576	0	0	23,788	23,788	-	並
	赤かび病	27,489	25.3	3,644	3.4	81,011	23,845	3,644	0	0	108,500	377,274	-	並
春まき小麦 (春まき) 16,828 ha	赤かび病	3,779	22.5	428	2.5	13,049	3,351	375	53	0	16,828	60,957	-	並
	ムギキモグリバエ	1,366	8.1	84	0.5	15,462	1,282	84	0	0	6,602	10,742	早	やや多
春まき小麦 (初冬まき) 772 ha	赤かび病	114	14.8	3	0.4	658	111	3	0	0	772	2,859	-	やや少
	ムギキモグリバエ	17	2.2	0	0.0	755	17	0	0	0	241	272	早	並

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積(ha)	率(%)	面積(ha)	率(%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
大豆 42,000 ha	べと病	8,985	21.4	13	0.0	33,015	8,972	13	0	0	19,507	23,690	並	並
	わい化病	3,320	7.9	0	0.0	38,680	3,320	0	0	0	42,000	61,372	-	やや少
	マメシクイガ	5,503	13.1	50	0.1	36,497	5,453	50	0	0	41,973	90,735	やや早	やや少
	食葉性鱗翅目幼虫	8,156	19.4	0	0.0	33,844	8,156	0	0	0	35,530	58,101	並	やや少
	タネバエ	180	0.4	8	0.0	41,820	172	8	0	0	42,000	42,000	-	並
小豆 19,000 ha	菌核病	1,471	7.7	2	0.0	17,529	1,469	2	0	0	18,904	56,723	並	やや少
	灰色かび病	1,930	10.2	0	0.0	17,070	1,930	0	0	0	18,935	56,490	並	やや少
	茎疫病	865	4.6	32	0.2	18,135	833	32	0	0	3,331	3,464	-	並
	落葉病	388	2.0	13	0.1	18,612	375	8	5	0	0	0	-	やや少
	食葉性鱗翅目幼虫	3,108	16.4	210	1.1	15,892	2,898	210	0	0	18,916	29,434	並	並
菜豆 6,660 ha	菌核病	1,264	19.0	16	0.2	5,396	1,248	16	0	0	6,445	19,793	並	並
	灰色かび病	1,296	19.5	16	0.2	5,364	1,280	16	0	0	6,445	19,793	やや遅	並
	黄化病	148	2.2	0	0.0	6,512	148	0	0	0	6,660	8,030	-	やや少
	タネバエ	49	0.7	0	0.0	6,611	49	0	0	0	6,660	6,660	-	並
ばれいしょ 47,100 ha	疫病	5,461	11.6	533	1.1	41,639	4,928	481	52	0	47,100	370,765	並	並
	塊茎腐敗	5,296	11.2	151	0.3	41,804	5,145	151	0	0	47,100	370,765	-	並
	軟腐病	8,081	17.2	644	1.4	39,019	7,437	627	17	0	46,900	159,447	-	並
	黒あし病	1,671	3.5	14	0.0	45,429	1,657	14	0	0	38,241	38,241	-	並
	そうか病	6,413	13.6	812	1.7	40,687	5,601	812	0	0	38,241	38,241	-	少
	粉状そうか病	3,949	8.4	185	0.4	43,151	3,764	185	0	0	3,129	3,129	-	並
	アブラムシ類	6,024	12.8	302	0.6	41,076	5,722	302	0	0	47,100	138,510	並	やや少
てんさい 57,700 ha	褐斑病	30,283	52.5	5,571	9.7	27,417	24,712	4,807	764	0	57,700	298,384	早	やや多
	根腐病 (黒根病を含む)	6,930	12.0	129	0.2	50,770	6,801	129	0	0	57,483	107,840	-	やや少
	そう根病	18	0.0	0	0.0	57,682	18	0	0	0	0	0	-	少
	黄化病 (西部萎黄病)	552	1.0	0	0.0	57,148	552	0	0	0	56,039	155,376	-	少
	ヨトウガ (第1回)	8,067	14.0	122	0.2	49,633	7,945	122	0	0	57,700	89,985	早	やや少
	ヨトウガ (第2回)	7,693	13.3	102	0.2	50,007	7,591	102	0	0	57,700	89,267	並	やや少
	テンサイ モグリハナバエ	32	0.1	0	0.0	57,668	32	0	0	0	49,285	63,882	並	並

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
たまねぎ 14,600 ha	白斑葉枯病	3,200	21.9	60	0.4	11,400	3,140	60	0	0	14,600	101,547	やや遅	少
	軟腐病	9,712	66.5	1,967	13.5	4,888	7,745	1,293	471	203	14,600	124,196	-	多
	乾腐病	863	5.9	5	0.0	13,737	858	5	0	0	0	0	-	少
	タマネギバエ、 タネバエ	720	4.9	22	0.2	13,880	698	22	0	0	6,251	7,114	-	並
	ネギアザミウマ	13,969	95.7	754	5.2	631	13,215	610	137	7	14,600	87,929	やや早	やや少
	ネギハモグリ バエ	1,852	12.7	44	0.3	12,748	1,808	40	4	0	13,726	55,859	やや早	やや少
ねぎ 654 ha	さび病	100	15.3	1	0.2	554	99	1	0	0	654	1,728	-	やや多
	ネギアザミウマ	489	74.8	51	7.8	165	438	50	1	0	654	3,918	並	やや少
にんじん 4,540 ha	黒葉枯病	1,542	34.0	211	4.6	2,998	1,331	140	71	0	4,383	15,537	-	やや多
だいこん 2,980 ha	軟腐病	948	31.8	180	6.0	2,032	768	180	0	0	2,818	8,488	-	並
	キスジ トビハムシ	719	24.1	10	0.3	2,261	709	10	0	0	2,980	6,290	-	やや少
キャベツ 1,180 ha	コナガ	411	34.8	6	0.5	769	405	6	0	0	1,180	6,863	早	少
	ヨトウガ	324	27.5	0	0.0	856	324	0	0	0	1,180	5,727	早	少
ブロッコリー 3,030 ha	コナガ	1,982	65.4	183	6.0	1,048	1,799	178	5	0	3,030	11,696	早	やや少
	ヨトウガ	1,407	46.4	48	1.6	1,623	1,359	45	3	0	3,030	9,455	早	やや少
りんご 518 ha	モニリア病	1	0.2	0	0.0	517	1	0	0	0	518	1,073	並	少
	黒星病	65	12.5	1	0.2	453	64	1	0	0	518	5,051	並	やや少
	斑点落葉病	80	15.4	3	0.6	438	77	3	0	0	518	2,530	並	並
	腐らん病	518	100.0	125	24.1	0	393	113	12	0	518	1,306	-	多
	ハマキムシ類	21	4.1	0	0.0	497	21	0	0	0	518	2,297	並	少
	モモシンクイガ	57	11.0	0	0.0	461	57	0	0	0	518	2,312	やや遅	やや少
	ハダニ類	110	21.2	16	3.1	408	94	13	3	0	518	1,546	やや遅	やや少
	キンモンホソガ	92	17.8	0	0.0	426	92	0	0	0	518	1,689	並	並
	ネズミ類	22	4.2	0	0.0	496	22	0	0	0	11	11	-	やや少