

V 主要病害虫発生程度別面積及び防除面積

作物名	病害虫名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
水稲 93,300 ha	葉いもち	256	0.3	79	0.1	93,044	177	31	48	0	91,199	218,554	並	並
	穂いもち	235	0.3	79	0.1	93,065	156	31	48	0	92,835	227,290	やや早	並
	紋枯病	18,210	19.5	1,218	1.3	75,090	16,992	1,184	34	0	18,966	21,943	並	並
	縞葉枯病	1,638	1.8	23	0.0	91,662	1,615	23	0	0	51,076	74,275	-	やや多
	ばか苗病	720	0.8	45	0.0	92,580	675	17	28	0	93,245	93,394	-	並
	苗立枯病	1,074	1.2	190	0.2	92,226	884	190	0	0	65,121	73,053	-	並
	種子伝染性細菌病	107	0.1	0	0.0	93,193	107	0	0	0	93,256	93,256	-	並
	ニカメイガ	777	0.8	0	0.0	92,523	777	0	0	0	43,738	76,653	並	並
	セジロウンカ	1,940	2.1	0	0.0	91,360	1,940	0	0	0	91,166	183,052	早	並
	ヒメトビウンカ	9,809	10.5	230	0.2	83,491	9,579	230	0	0	87,672	183,312	やや早	やや少
	イネドロオイムシ	2,154	2.3	21	0.0	91,146	2,133	21	0	0	90,056	91,993	並	少
	アカヒゲ ホソミドリ カスミカメ	29,462	31.6	1,139	1.2	63,838	28,323	1,139	0	0	93,190	219,231	やや早	やや多
	イネミギワバエ	248	0.3	0	0.0	93,052	248	0	0	0	27,185	27,185	並	少
フタオビコヤガ	1,049	1.1	0	0.0	92,251	1,049	0	0	0	79,699	104,693	並	少	
秋まき小麦 132,300 ha	赤さび病	59,055	44.6	17,081	12.9	73,245	41,974	14,641	2,108	332	132,152	357,854	早	多
	うどんこ病	10,770	8.1	34	0.0	121,530	10,736	34	0	0	132,152	321,985	並	並
	雪腐病類	19,458	14.7	1,926	1.5	112,842	17,532	1,703	191	32	128,184	131,524	-	少
	眼紋病	8,159	6.2	1,527	1.2	124,141	6,632	1,489	38	0	32,297	32,297	-	やや多
	赤かび病	8,163	6.2	0	0.0	124,137	8,163	0	0	0	132,300	448,792	-	少
春まき小麦 (春まき) 17,660 ha	赤かび病	1,092	6.2	53	0.3	16,568	1,039	53	0	0	17,660	62,617	-	少
	ムギキモグリバエ	1,000	5.7	0	0.0	16,660	1,000	0	0	0	6,954	11,655	早	並
春まき小麦 (初冬まき) 824 ha	赤かび病	20	2.4	0	0.0	804	20	0	0	0	824	2,681	-	少
	ムギキモグリバエ	95	11.5	0	0.0	729	95	0	0	0	558	558	早	多

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
大豆 45,300 ha	べと病	8,057	17.8	0	0.0	37,243	8,057	0	0	0	22,119	25,880	並	やや少
	わい化病	7,142	15.8	183	0.4	38,158	6,959	175	8	0	45,300	77,637	-	やや多
	マメシンクイガ	8,340	18.4	0	0.0	36,960	8,340	0	0	0	45,285	97,198	やや早	やや少
	食葉性 鱗翅目幼虫	10,898	24.1	283	0.6	34,402	10,615	258	25	0	40,602	71,142	やや早	並
	タネバエ	1,372	3.0	68	0.2	43,928	1,304	34	34	0	44,051	44,051	-	多
小豆 20,900 ha	菌核病	1,128	5.4	3	0.0	19,772	1,125	3	0	0	20,490	63,699	並	やや少
	灰色かび病	1,937	9.3	8	0.0	18,963	1,929	8	0	0	20,856	63,885	並	やや少
	茎疫病	248	1.2	41	0.2	20,652	207	41	0	0	3,348	4,583	-	やや少
	落葉病	340	1.6	0	0.0	20,560	340	0	0	0	0	0	-	並
	食葉性 鱗翅目幼虫	4,107	19.7	186	0.9	16,793	3,921	186	0	0	20,900	31,066	やや早	並
菜豆 5,450 ha	菌核病	274	5.0	0	0.0	5,176	274	0	0	0	5,097	15,457	やや遅	少
	灰色かび病	458	8.4	34	0.6	4,992	424	34	0	0	5,450	16,338	並	やや少
	黄化病	85	1.6	0	0.0	5,365	85	0	0	0	5,450	7,558	-	並
	タネバエ	13	0.2	0	0.0	5,437	13	0	0	0	5,450	0	-	少
ばれいしょ 48,500 ha	疫病	1,578	3.3	0	0.0	46,922	1,578	0	0	0	48,500	389,909	やや遅	少
	塊茎腐敗	2,097	4.3	0	0.0	46,403	2,097	0	0	0	48,500	389,909	-	少
	軟腐病	8,547	17.6	970	2.0	39,953	7,577	820	150	0	48,500	179,810	-	並
	黒あし病	1,163	2.4	15	0.0	47,337	1,148	0	15	0	38,475	38,475	-	並
	そうか病	13,787	28.4	3,641	7.5	34,713	10,146	2,656	806	179	38,563	38,563	-	やや多
	粉状そうか病	1,646	3.4	0	0.0	46,854	1,646	0	0	0	3,013	3,013	-	少
	アブラムシ類	6,202	12.8	9	0.0	42,298	6,193	9	0	0	48,500	145,451	早	少
てんさい 51,200 ha	褐斑病	44,511	86.9	15,975	31.2	6,689	28,536	13,068	2,497	410	51,200	355,723	早	多
	根腐病 (黒根病を含む)	2,713	5.3	54	0.1	48,487	2,659	54	0	0	50,999	97,024	-	少
	そう根病	10	0.0	0	0.0	51,190	10	0	0	0	0	0	-	少
	黄化病 (西部萎黄病)	752	1.5	0	0.0	50,448	752	0	0	0	49,932	128,515	-	少
	ヨトウガ (第1回)	10,107	19.7	441	0.9	41,093	9,666	441	0	0	51,200	86,837	並	並
	ヨトウガ (第2回)	9,206	18.0	374	0.7	41,994	8,832	374	0	0	51,200	81,736	並	やや多
	テンサイ モグリハナバエ	468	0.9	0	0.0	50,732	468	0	0	0	49,642	77,333	並	並

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
たまねぎ 14,900 ha	白斑葉枯病	2,367	15.9	8	0.1	12,533	2,359	8	0	0	14,900	103,855	並	少
	軟腐病	3,868	26.0	18	0.1	11,032	3,850	18	0	0	14,900	116,928	-	少
	乾腐病	1,341	9.0	0	0.0	13,559	1,341	0	0	0	0	0	-	少
	タマネギバエ、 タネバエ	575	3.9	1	0.0	14,325	574	1	0	0	6,020	6,437	-	少
	ネギアザミウマ	13,985	93.9	639	4.3	915	13,346	531	108	0	14,900	110,648	早	やや少
	ネギハモグリ バエ	1,978	13.3	59	0.4	12,922	1,919	59	0	0	14,357	71,136	やや早	やや少
ねぎ 576 ha	さび病	118	20.5	0	0.0	458	118	0	0	0	552	1,976	-	並
	ネギアザミウマ	413	71.7	98	17.0	163	315	89	9	0	576	4,383	並	並
	ネギコガ	0	0.0	0	0.0	576	0	0	0	0	545	1,731	-	-
	ネギハモグリ バエ	39	6.8	0	0.0	537	39	0	0	0	545	2,256	-	-
ながいも 1,790 ha	アブラムシ類	11	0.6	0	0.0	1,779	11	0	0	0	688	3,626	-	-
にんじん 4,281 ha	黒葉枯病	417	9.7	16	0.4	3,864	401	16	0	0	4,088	13,166	-	少
だいこん 2,632 ha	軟腐病	827	31.4	135	5.1	1,805	692	87	48	0	2,385	7,680	-	並
	キスジ トビハムシ	613	23.3	10	0.4	2,019	603	10	0	0	2,632	8,101	-	やや少
キャベツ 1,044 ha	コナガ	562	53.8	25	2.4	482	537	25	0	0	1,044	6,681	早	やや少
	ヨトウガ	229	21.9	4	0.4	815	225	4	0	0	1,044	5,135	並	少
ブロッコリー 3,230 ha	コナガ	2,324	72.0	170	5.3	906	2,154	170	0	0	3,230	14,002	早	少
	ヨトウガ	1,135	35.1	45	1.4	2,095	1,090	45	0	0	3,230	10,784	並	少
りんご 512 ha	モニリア病	3	0.6	0	0.0	509	3	0	0	0	512	1,060	並	並
	黒星病	10	2.0	0	0.0	502	10	0	0	0	512	4,991	並	少
	斑点落葉病	39	7.6	0	0.0	473	39	0	0	0	512	2,534	やや早	少
	腐らん病	512	100.0	95	18.6	0	417	82	13	0	512	1,328	-	多
	ハマキムシ類	20	3.9	0	0.0	492	20	0	0	0	512	2,343	早	少
	モモシクイガ	43	8.4	1	0.2	469	42	1	0	0	512	2,357	やや早	並
	ハダニ類	29	5.7	3	0.6	483	26	3	0	0	512	1,520	並	少
	キンモンホソガ	69	13.5	0	0.0	443	69	0	0	0	512	1,668	並	やや少