

V 主要病害虫発生程度別面積及び防除面積

作物名	病害虫名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
水稲 95,000 ha	葉いもち	446	0.5	0	0.0	94,554	446	0	0	0	94,560	221,087	並	並
	穂いもち	602	0.6	16	0.0	94,398	586	16	0	0	94,560	221,087	並	並
	紋枯病	19,968	21.0	1,718	1.8	75,032	18,250	1,502	216	0	17,362	20,252	やや早	並
	縞葉枯病	1,440	1.5	0	0.0	93,560	1,440	0	0	0	50,861	63,238	-	並
	ばか苗病	550	0.6	43	0.0	94,450	507	43	0	0	94,957	94,957	-	並
	苗立枯病	1,328	1.4	50	0.1	93,672	1,278	50	0	0	76,918	80,665	-	並
	種子伝染性細菌病	60	0.1	43	0.0	94,940	17	43	0	0	94,957	94,957	-	並
	ニカメイガ	727	0.8	0	0.0	94,273	727	0	0	0	50,679	83,239	並	並
	セジロウンカ	1,227	1.3	0	0.0	93,773	1,227	0	0	0	93,015	193,059	やや早	並
	ヒメトビウンカ	21,194	22.3	9	0.0	73,806	21,185	9	0	0	89,092	183,167	やや遅	やや多
	イネドロオイムシ	2,919	3.1	209	0.2	92,081	2,710	209	0	0	88,863	95,715	並	少
	アカヒゲ ホソミドリ カスミカメ	33,029	34.8	2,935	3.1	61,971	30,094	2,866	69	0	94,974	228,288	早	多
	イネミギワバエ	614	0.6	0	0.0	94,386	614	0	0	0	31,155	31,155	並	少
フタオビコヤガ	627	0.7	0	0.0	94,373	627	0	0	0	85,481	120,381	並	少	
秋まき小麦 132,300 ha	赤さび病	31,619	23.9	5,811	4.4	100,681	25,808	4,928	830	53	132,300	395,288	早	並
	うどんこ病	6,435	4.9	105	0.1	125,865	6,330	105	0	0	132,300	366,069	並	並
	雪腐病類	18,193	13.8	1,336	1.0	114,107	16,857	1,205	110	21	130,809	130,808	-	少
	眼紋病	7,612	5.8	1,276	1.0	124,688	6,336	1,254	22	0	34,940	34,941	-	やや多
	赤かび病	8,146	6.2	0	0.0	124,154	8,146	0	0	0	132,300	448,308	-	少
春まき小麦 (春まき) 17,797 ha	赤かび病	1,415	8.0	0	0.0	16,382	1,415	0	0	0	17,797	59,200	-	少
	ムギキモグリバエ	1,013	5.7	0	0.0	16,784	1,013	0	0	0	7,176	12,157	並	並
春まき小麦 (初冬まき) 868 ha	赤かび病	50	5.8	0	0.0	818	50	0	0	0	868	3,122	-	少
	ムギキモグリバエ	46	5.3	0	0.0	822	46	0	0	0	290	490	並	やや多

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
大豆 46,800 ha	べと病	7,579	16.2	0	0.0	39,221	7,579	0	0	0	21,154	24,973	並	やや少
	わい化病	6,136	13.1	164	0.4	40,664	5,972	82	82	0	46,787	82,314	-	やや多
	マメシンクイガ	8,701	18.6	95	0.2	38,099	8,606	95	0	0	46,783	102,042	早	やや少
	食葉性 鱗翅目幼虫	12,601	26.9	442	0.9	34,199	12,159	388	54	0	42,000	83,044	やや早	並
	タネバエ	311	0.7	0	0.0	46,489	311	0	0	0	46,787	46,787	-	並
小豆 20,800 ha	菌核病	1,132	5.4	0	0.0	19,668	1,132	0	0	0	20,415	66,353	並	やや少
	灰色かび病	2,871	13.8	0	0.0	17,929	2,871	0	0	0	20,800	66,572	並	並
	茎疫病	321	1.5	0	0.0	20,479	321	0	0	0	3,140	3,971	-	やや少
	落葉病	484	2.3	3	0.0	20,316	481	3	0	0	0	0	-	並
	食葉性 鱗翅目幼虫	4,450	21.4	66	0.3	16,350	4,384	66	0	0	20,602	38,496	並	並
菜豆 5,270 ha	菌核病	292	5.5	0	0.0	4,978	292	0	0	0	4,944	15,505	並	少
	灰色かび病	341	6.5	0	0.0	4,929	341	0	0	0	5,270	15,832	並	やや少
	黄化病	47	0.9	0	0.0	5,223	47	0	0	0	5,270	6,104	-	少
	タネバエ	27	0.5	0	0.0	5,243	27	0	0	0	5,270	0	-	並
ばれいしょ 48,700 ha	疫病	1,102	2.3	57	0.1	47,598	1,045	57	0	0	48,700	379,088	遅	少
	塊茎腐敗	1,758	3.6	35	0.1	46,942	1,723	35	0	0	48,700	379,088	-	少
	軟腐病	8,217	16.9	463	1.0	40,483	7,754	463	0	0	48,700	178,540	-	並
	黒あし病	1,413	2.9	2	0.0	47,287	1,411	2	0	0	38,453	38,792	-	並
	そうか病	12,417	25.5	2,545	5.2	36,283	9,872	1,814	561	170	38,453	38,453	-	並
	粉状そうか病	2,292	4.7	69	0.1	46,408	2,223	69	0	0	3,928	3,928	-	やや少
	アブラムシ類	5,689	11.7	333	0.7	43,011	5,356	333	0	0	48,700	146,568	やや早	やや少
てんさい 48,900 ha	褐斑病	39,435	80.6	8,335	17.0	9,465	31,100	6,426	1,762	147	48,900	346,041	早	多
	根腐病 (黒根病を含む)	2,932	6.0	0	0.0	45,968	2,932	0	0	0	48,701	97,691	-	少
	そう根病	52	0.1	0	0.0	48,848	52	0	0	0	0	0	-	やや少
	黄化病 (西部萎黄病)	570	1.2	0	0.0	48,330	570	0	0	0	47,470	122,456	-	少
	ヨトウガ (第1回)	9,533	19.5	505	1.0	39,367	9,028	496	9	0	48,900	84,665	やや早	やや多
	ヨトウガ (第2回)	17,731	36.3	4,470	9.1	31,169	13,261	4,313	135	22	48,900	89,292	やや早	多
	テンサイ モグリハナバエ	1,505	3.1	288	0.6	47,395	1,217	288	0	0	40,463	54,608	並	やや多

作物名	病虫害名	発生面積		被害面積		発生程度別面積 (ha)					防除面積 (ha)		概評	
		面積 (ha)	率 (%)	面積 (ha)	率 (%)	無	少	中	多	甚	実面積	延面積	初発期	発生量
たまねぎ 14,900 ha	白斑葉枯病	698	4.7	0	0.0	14,202	698	0	0	0	14,900	102,194	遅	少
	軟腐病	3,836	25.7	40	0.3	11,064	3,796	40	0	0	14,900	112,451	-	少
	乾腐病	1,525	10.2	70	0.5	13,375	1,455	70	0	0	151	151	-	少
	タマネギバエ、 タネバエ	444	3.0	18	0.1	14,456	426	18	0	0	5,840	6,259	-	少
	ネギアザミウマ	14,048	94.3	1,730	11.6	852	12,318	1,480	250	0	14,900	110,352	早	やや多
	ネギハモグリ バエ	702	4.7	39	0.3	14,198	663	39	0	0	13,288	81,274	並	少
ねぎ 602 ha	さび病	37	6.1	0	0.0	565	37	0	0	0	577	2,073	-	少
	ネギアザミウマ	576	95.7	304	50.5	26	272	249	55	0	602	4,611	やや早	多
	ネギコガ	2	0.3	0	0.0	600	2	0	0	0	602	1,890	-	-
	ネギハモグリ バエ	63	10.5	7	1.2	539	56	7	0	0	602	2,368	-	-
ながいも 1,720 ha	アブラムシ類	22	1.3	0	0.0	1,698	22	0	0	0	808	4,148	-	-
にんじん 4,190 ha	黒葉枯病	401	9.6	8	0.2	3,789	393	8	0	0	4,003	13,378	-	少
だいこん 2,600 ha	軟腐病	835	32.1	146	5.6	1,765	689	95	51	0	2,366	8,364	-	並
	キスジ トビハムシ	584	22.5	22	0.8	2,016	562	22	0	0	2,600	8,056	-	やや少
キャベツ 1,060 ha	コナガ	588	55.5	54	5.1	472	534	54	0	0	1,060	6,797	早	やや多
	ヨトウガ	240	22.6	15	1.4	820	225	15	0	0	1,060	5,248	やや早	やや少
ブロッコリー 3,290 ha	コナガ	2,324	70.6	280	8.5	966	2,044	255	25	0	3,290	14,352	早	並
	ヨトウガ	1,105	33.6	10	0.3	2,185	1,095	10	0	0	3,290	11,126	やや早	少
りんご 501 ha	モニリア病	15	3.0	0	0.0	486	15	0	0	0	501	1,002	並	並
	黒星病	4	0.8	0	0.0	497	4	0	0	0	501	5,006	並	少
	斑点落葉病	36	7.2	0	0.0	465	36	0	0	0	501	2,657	並	少
	腐らん病	502	100.2	117	23.4	0	385	104	13	0	501	1,495	-	多
	ハマキムシ類	27	5.4	0	0.0	474	27	0	0	0	501	2,204	やや遅	少
	モモシクイガ	72	14.4	2	0.4	429	70	1	1	0	501	1,897	早	並
	ハダニ類	15	3.0	1	0.2	486	14	1	0	0	501	1,161	並	少
	キンモンホソガ	69	13.8	0	0.0	432	69	0	0	0	501	1,846	やや早	やや少